

COMPLETE GRADIENT SHRINKING RICCI SOLITONS HAVE FINITE TOPOLOGICAL TYPE

FUQUAN FANG, JIANWEN MAN, AND ZHENLEI ZHANG

ABSTRACT. We show that a complete Riemannian manifold has finite topological type (i.e., homeomorphic to the interior of a compact manifold with boundary), provided its Bakry-Émery Ricci tensor has a positive lower bound, and either of the following conditions:

- (i) the Ricci curvature is bounded from above;
- (ii) the Ricci curvature is bounded from below and injectivity radius is bounded away from zero.

Moreover, a complete shrinking Ricci soliton has finite topological type if its scalar curvature is bounded.

1. INTRODUCTION

In 1968, J. Milnor [8] conjectured that a complete non-compact Riemannian manifold with non-negative Ricci curvature has a finitely generated fundamental group. However, such a manifold may not have finite topological type. Examples of complete non-compact manifold with positive Ricci curvature without finite topological type was constructed by Gromoll-Meyer [5]. It has been an interesting topic in Riemannian geometry to study the topology of complete manifolds with positive (non-negative) Ricci curvature.

In this note we are concerned with complete Riemannian manifold (M, g) satisfying that $\text{Ric} + \text{Hess}(f) \geq \lambda g$ for some constant $\lambda > 0$ and $f \in C^\infty(M)$, i.e., whose Bakry-Émery Ricci tensor is bounded below by λ in the sense of [7]. When the equality holds, the manifold is a shrinking Ricci soliton, i.e., a self-similar solution of the well-known Ricci flow equation. If f is constant, Bakry-Émery Ricci tensor reduces to the Ricci tensor, and so the classical Myers' theorem implies that M is compact with finite fundamental group. In general, M may not be compact, but from the work of [4, 6, 7, 9, 10, 11] etc., M still has finite fundamental group.

The main result of this note shows that a complete Riemannian manifold whose Bakry-Émery Ricci tensor is bounded below by $\lambda > 0$ has finite topological type, provided the Ricci curvature is bounded from above. Moreover,

2000 Mathematics Subject Classification. Primary 53C25; Secondary 53C21, 55Q52 .

Key words and phrases. Ricci soliton, finite topological type.

Supported by NSF Grant of China #10671097 and the Capital Normal University.

a shrinking Ricci soliton has finite topological type if its scalar curvature is bounded.

Theorem 1. *Suppose (M, g) is a complete Riemannian manifold satisfying $\text{Ric} + \text{Hess}(f) \geq \lambda g$ for some constant $\lambda > 0$ and $f \in C^\infty(M)$. Then M is of finite topological type, if either of the following alternative conditions holds:*

- (i) $\text{Ric} \leq Cg$ for some constant $C < \infty$;
- (ii) $\text{Ric} \geq -\delta^{-1}g$ and the injectivity radius $\text{inj}(M, g) \geq \delta > 0$ for some $\delta > 0$.

If (M, g) is a shrinking Ricci soliton, then the Ricci curvature bounds can be relaxed by scalar curvature.

Theorem 2. *Suppose (M, g) is a complete shrinking Ricci soliton $\text{Ric} + \text{Hess}(f) = \frac{g}{2}$, where $f \in C^\infty(M)$. If the scalar curvature R is bounded, then M has finite topological type.*

In view of Theorem 2 it is nature to pose the following

Conjecture 3. *Any shrinking Ricci soliton has finite topological type.*

We prove Theorem 1 in section 2 and Theorem 2 in section 3.

2. PROOF OF THEOREM 1

Let (M, g) be such a manifold satisfying that $\text{Ric} + \text{Hess}(f) \geq \lambda g$ for some $\lambda > 0$ and $f \in C^\infty(M)$. By the deformation lemma of Morse theory, to prove Theorem 1, it suffices to show that the function f is proper and has no critical points outside of a compact set.

First fix one point $p \in M$ as a base point. For any $q \in M$ with $d(p, q) = L$, choose a shortest geodesic γ from p to q parametrized by arc length. Then

$$\begin{aligned} \langle \nabla f, \dot{\gamma} \rangle(q) &= \langle \nabla f, \dot{\gamma} \rangle(p) + \int_0^L \frac{d^2}{dt^2} f(\gamma(t)) dt \\ &\geq \langle \nabla f, \dot{\gamma} \rangle(p) + \int_0^L (\lambda - \text{Ric}(\dot{\gamma}, \dot{\gamma})) dt \\ &\geq \lambda L - |\nabla f|(p) - \int_0^L \text{Ric}(\dot{\gamma}, \dot{\gamma}) dt. \end{aligned}$$

If the integral

$$\int_0^L \text{Ric}(\dot{\gamma}, \dot{\gamma}) dt \leq \Lambda$$

for some constant Λ independent of q and the choice of γ , then

$$|\nabla f|(q) \geq \langle \nabla f, \dot{\gamma} \rangle(q) \geq \lambda d(p, q) - |\nabla f|(p) - \Lambda,$$

which implies that $|\nabla f|(q)$ has a linear growth in $d(p, q)$ and so f is a proper function without critical points outside of a compact set. In the remainder of this section, we will focus on proving that $\int_0^L Ric(\dot{\gamma}, \dot{\gamma})$ has an upper bound under the assumptions of Theorem 1.

Case (i): $Ric \leq Cg$ for some constant $C < \infty$;

By Lemma 2.2 of [9], the integral bound is given by $\Lambda = 2(n-1) + 2C$.

Case (ii): $Ric \geq -\delta^{-1}g$ and $\text{inj}(M, g) \geq \delta > 0$ for some $\delta > 0$.

Suppose $d(p, q) = L \geq \delta$. Let $\varphi(t) : [0, L] \rightarrow [0, 1]$ be an arcwise smooth function such that $\varphi(0) = \varphi(L) = 0$. By the second variation formula, as did in [9] or [11], we have the following estimate:

$$\int_0^L \varphi^2(t) Ric(\dot{\gamma}, \dot{\gamma}) dt \leq (n-1) \int_0^L |\dot{\varphi}|^2 dt.$$

Now define φ by

$$\varphi = \begin{cases} \frac{3}{\delta}t, & t \in [0, \frac{\delta}{3}]; \\ 1, & t \in [\frac{\delta}{3}, L - \frac{\delta}{3}]; \\ \frac{3}{\delta}(L-t), & t \in [L - \frac{\delta}{3}, L], \end{cases}$$

then we have the estimate

$$\begin{aligned} \int_0^L Ric(\dot{\gamma}, \dot{\gamma}) dt &\leq (n-1) \int_0^L |\dot{\varphi}|^2 dt + \int_0^{\frac{\delta}{3}} (1 - \varphi^2) Ric(\dot{\gamma}, \dot{\gamma}) dt + \int_{L-\frac{\delta}{3}}^L (1 - \varphi^2) Ric(\dot{\gamma}, \dot{\gamma}) dt \\ &\leq \frac{6}{\delta}(n-1) + \frac{2}{3} + \int_0^{\frac{\delta}{3}} Ric(\dot{\gamma}, \dot{\gamma}) dt + \int_{L-\frac{\delta}{3}}^L Ric(\dot{\gamma}, \dot{\gamma}) dt, \end{aligned}$$

where in the second inequality, we used the fact that

$$\begin{aligned} \int_0^{\frac{\delta}{3}} (1 - \varphi^2) Ric(\dot{\gamma}, \dot{\gamma}) dt &= \int_0^{\frac{\delta}{3}} (1 - \varphi^2) (Ric(\dot{\gamma}, \dot{\gamma}) + \frac{1}{\delta}) dt - \frac{1}{\delta} \int_0^{\frac{\delta}{3}} (1 - \varphi^2) dt \\ &\leq \int_0^{\frac{\delta}{3}} (Ric(\dot{\gamma}, \dot{\gamma}) + \frac{1}{\delta}) dt - \frac{1}{\delta} \int_0^{\frac{\delta}{3}} (1 - \varphi^2) dt \\ &\leq \int_0^{\frac{\delta}{3}} Ric(\dot{\gamma}, \dot{\gamma}) dt + \frac{1}{\delta} \int_0^{\frac{\delta}{3}} \varphi^2 dt \\ &\leq \int_0^{\frac{\delta}{3}} Ric(\dot{\gamma}, \dot{\gamma}) dt + \frac{1}{3}, \end{aligned}$$

and similarly

$$\int_{L-\frac{\delta}{3}}^L (1 - \varphi^2) Ric(\dot{\gamma}, \dot{\gamma}) dt \leq \int_{L-\frac{\delta}{3}}^L Ric(\dot{\gamma}, \dot{\gamma}) dt + \frac{\delta}{3}.$$

We next prove that $\int_0^{\frac{\delta}{3}} Ric(\dot{\gamma}, \dot{\gamma})dt$ and $\int_{L-\frac{\delta}{3}}^L Ric(\dot{\gamma}, \dot{\gamma})dt$ are bounded from above and so finish the proof of Theorem 1. This is given by the following lemma.

Lemma 4. *If $Ric \geq -\delta^{-1}g$ and $\text{inj}(M, g) \geq \delta > 0$ for some $\delta > 0$, then*

$$\int_0^{\frac{\delta}{3}} Ric(\dot{\gamma}, \dot{\gamma})dt \leq \frac{6}{\delta}(n-1) + \frac{2}{3}$$

for any minimal arc length parametrized geodesic $\gamma : [0, \frac{\delta}{3}] \rightarrow M$.

Proof. Firstly, by $\text{inj}(M, g) \geq \delta$, we can extend the geodesic γ to a shortest geodesic $\sigma : [0, \delta] \rightarrow M$, such that $\gamma(t) = \sigma(t + \frac{\delta}{3}), t \in [0, \frac{\delta}{3}]$.

Set $L = \delta$ in the arguments above, we have

$$\int_0^{\delta} \varphi^2 Ric(\dot{\sigma}, \dot{\sigma})dt \leq (n-1) \int_0^{\delta} |\dot{\sigma}|^2 dt = \frac{6}{\delta}(n-1),$$

then using $Ric \geq -\delta^{-1}g$, we get the estimate

$$\begin{aligned} \int_0^{\frac{\delta}{3}} Ric(\dot{\gamma}, \dot{\gamma})dt &= \int_{\frac{\delta}{3}}^{\frac{2\delta}{3}} Ric(\dot{\sigma}, \dot{\sigma})dt \\ &\leq \frac{6}{\delta}(n-1) - \int_0^{\frac{\delta}{3}} \varphi^2 Ric(\dot{\sigma}, \dot{\sigma})dt - \int_{\frac{2\delta}{3}}^{\delta} \varphi^2 Ric(\dot{\sigma}, \dot{\sigma})dt \\ &\leq \frac{6}{\delta}(n-1) + \frac{2}{3}. \end{aligned}$$

This concludes the result. \square

3. PROOF OF THEOREM 2

As before, we will prove that the potential function f to the Ricci soliton is proper and has no critical points outside of $B(p, \rho)$ for large ρ .

Suppose (M, g) is a complete shrinking Ricci soliton which satisfies

$$(1) \quad Ric + \text{Hess}(f) = \frac{g}{2}$$

for some potential function f . Suppose further that the scalar curvature $|R| \leq C$ for some constant $C < \infty$. It's well-known that the following analytic equality holds for the soliton (after modifying f by a translation, see [3] for example.):

$$(2) \quad R + |\nabla f|^2 = f.$$

We begin with several lemmas. Let $q \in M$ be one critical point of f and denote by $\rho = d(p, q)$ the distance from p to q . Let γ be a shortest arc length parametrized geodesic from p to q . Then we have

Lemma 5.

$$(3) \quad \frac{\rho}{2} - |\nabla f|(p) \leq \int_0^\rho Ric(\dot{\gamma}, \dot{\gamma}) dt.$$

Proof. By a direct computation,

$$\begin{aligned} 0 = \langle \nabla f, \dot{\gamma} \rangle(q) &= \langle \nabla f, \dot{\gamma} \rangle(p) + \int_0^\rho \frac{d^2}{dt^2} f(\gamma(t)) dt \\ &\geq -|\nabla f|(p) + \int_0^\rho \left(\frac{1}{2} - Ric(\dot{\gamma}, \dot{\gamma}) \right) dt. \end{aligned}$$

Then the result follows. \square

On the other hand, by second variation formula as did in above section, we can get an upper bound for $\int_0^\rho Ric(\dot{\gamma}, \dot{\gamma}) dt$. Precisely, for the function ψ defined by

$$\psi(t) = t, t \in [0, 1]; \psi(t) \equiv 1, t \in [1, \rho_i - 1]; \psi(t) = \rho_i - t, t \in [\rho_i - 1, \rho_i],$$

we have the estimate

$$\begin{aligned} \int_0^\rho Ric(\dot{\gamma}, \dot{\gamma}) dt &\leq \int_0^\rho (n-1) |\dot{\psi}|^2 dt + \int_0^1 (1-\psi^2) Ric(\dot{\gamma}, \dot{\gamma}) dt \\ &\quad + \int_{\rho-1}^\rho (1-\psi^2) Ric(\dot{\gamma}, \dot{\gamma}) dt \\ &\leq 2(n-1) + \sup_{B(p,1)} |Ric| + \int_{\rho-1}^\rho (1-\psi^2) \left(\frac{1}{2} - \frac{d^2}{dt^2} f(\gamma(t)) \right) dt \\ &\leq 2(n-1) + 1 + \sup_{B(p,1)} |Ric| - \int_{\rho-1}^\rho (1-\psi^2) \frac{d^2}{dt^2} f(\gamma(t)) dt. \end{aligned}$$

Do integration by parts, we have the estimate for the last term

$$\begin{aligned} - \int_{\rho-1}^\rho (1-\psi^2) \frac{d^2}{dt^2} f(\gamma(t)) dt &= 2 \int_{\rho-1}^\rho \psi \frac{d}{dt} f(\gamma(t)) dt \\ &= -2f(\gamma(\rho-1)) + 2 \int_{\rho-1}^\rho f(\gamma(t)) dt. \end{aligned}$$

Substituting this equality into above estimate, we obtain

Lemma 6.

$$\begin{aligned} \int_0^\rho Ric(\dot{\gamma}, \dot{\gamma}) dt &\leq 2n + \sup_{B(p,1)} |Ric| + 2 \int_{\rho-1}^\rho f(\gamma(t)) dt - 2f(\gamma(\rho-1)) \\ &\leq 2n + \sup_{B(p,1)} |Ric| + \sup_{x,y \in B(q,1)} 2|f(x) - f(y)|. \end{aligned}$$

Now we use equation (2) to give an upper bound of $\int_0^\rho Ric(\dot{\gamma}, \dot{\gamma})dt$. First by equation (2) we have the gradient estimate $|\nabla f| \leq \sqrt{f-R} \leq \sqrt{f+C}$. Then by assumption, q is a critical point of f , so $|f(q)| = |R(q)| \leq C$. Integrating along a geodesic, we see that for any $x \in B(q, 1)$

$$(4) \quad \sqrt{f(x) + C} \leq \sqrt{f(q) + C} + \frac{d(x, q)}{2} \leq \sqrt{2C} + 1.$$

Thus $f(x) \leq 3C + 2$ for all $x \in B(q, 1)$ and consequently

$$(5) \quad \sup_{x, y \in B(q, 1)} |f(x) - f(y)| \leq (3C + 2 + C) = 4C + 2.$$

The combination of Lemma 5, Lemma 6 and equation (5) gives the upper bound of the distance $\rho = d(p, q)$:

$$\rho \leq 4n + 8C + 4 + 2|\nabla f|(p) + \sup_{B(p, 1)} 2|Ric|.$$

Note that the arguments above just used the upper boundedness of f . By the same reason as before, we conclude that f is proper and then finish the proof of the theorem.

REFERENCES

- [1] W. Ambrose, *A theorem of Myers*, Duke Math. J. 24 (1957), 345-348.
- [2] A. Derdzinski, *A Myers-type theorem and compact Ricci solitons*, Proc. AMS., 134 (2006), 3645-3648.
- [3] H.D. Cao, R. Hamilton and T. Ilmanen, *Gauss densities and stability for some Ricci solitons*, arXiv:math.DG/0404165.
- [4] M. Fernández-López and E. García-Río, *A remark on compact Ricci solitons*, Math. Ann., DOI 10.1007/s00208-007-0173-4.
- [5] D. Gromoll and W. T. Meyer, *Examples of complete manifolds with positive Ricci curvature*, J. Differential Geom. 21 (1985), no. 2, 195-211.
- [6] X.M. Li, *On extensions of Myers' theorem*, Bull. London Math. Soc., 27 (1995), 392-396.
- [7] J. Lott, *Some geometric properties of the Bakry-Émery-Ricci tensor*, Comment. Math. Helv. 78 (2003), 865-883.
- [8] J. Milnor, *Growth of finitely generated solvable groups*, J. Differential Geometry 2(1968), 447-449.
- [9] W. Wylie, *Complete shrinking Ricci solitons have finite fundamental group*, to appear in Proc. AMS.
- [10] G.F. Wei and W. Wylie, *Comparison geometry for the Bakry-Émery Ricci tensor*, arXiv:math.DG/0706.1120v1.
- [11] Z.L. Zhang, *On the finiteness of the fundamental group of a compact shrinking Ricci soliton*, Collq. Math., 107 (2007), 297-299.

DEPARTMENT OF MATHEMATICS, CAPITAL NORMAL UNIVERSITY, BEIJING, P.R.
CHINA

E-mail address: fuquan_fang@yahoo.com

NANKAI INSTITUTE OF MATHEMATICS, WEIJIN ROAD 94, TIANJIN 300071, P.R.
CHINA

NANKAI INSTITUTE OF MATHEMATICS, WEIJIN ROAD 94, TIANJIN 300071, P.R.
CHINA

E-mail address: zhleigo@yahoo.com.cn